



# Pressemitteilung

## How Touch and Movement Contribute to the Development of the Brain

**Neuroscientist at the Excellence Cluster CIN at the University of Tübingen together with French colleagues uncovered in an animal model the neuronal processes that underlay the development of sensory maps in the developing brain.**

Tübingen, 14.10.2011

Every expectant mother is aware of fetal movements in the late stages of pregnancy. It is known that the frequency of fetal movements is correlated with the physical fitness of the newborn child. What is the functional role of these irregular, non-coordinated movements in the brain development? And what are the neuronal processes that facilitate the brain development in result of these movements? The Neuroscientists Dr. Anton Sirota from the Excellence Cluster Werner Reichardt-Center for Integrative Neuroscience (CIN) at the University of Tübingen and Dr. Rustem Khazipov from the Institut National de la Santé et de la Recherche Médicale (INSERM) in France pursue these questions in an intensive and long standing collaboration. In an article published in the current issue of the scientific journal *Science* they could show that this process is controlled by so called early gamma oscillations (EGO) in the developing brain.

In the first week of life newborn rats are at a similar developmental stage as children in the third trimester of pregnancy. Newborn rats display perpetual twitches and jerks reminiscent of the human fetal movements. These spontaneous twitches as well as passive touches help to establish neuronal topographical maps of the body parts in the brain. Each stimulation of a single whisker (through twitches of the snout or the touch of mother or littermates) results in a unique pattern of neural activity, that the authors termed "early gamma oscillations" (EGO), which are exclusively confined to neural circuits of the thalamus and neocortex, which are genetically pre-wired to represent this particular whisker. The sensory information of the whisker and the neuronal activity during development are instrumental for establishing a functional topographic map of the sensory information. The high frequency of EGO of about 40 Hz is essential for strengthening neuronal connections. Every repetition strengthens further the connections between neurons in cortex and thalamus into a topographic and functional unit. During the maturation of the brain and the neuronal machinery, the EGO gradually disappear and they are being replaced by gamma oscillations of the adult which serve horizontal binding and other integrative cortical functions in the mature brain.

### Hochschulkommunikation

**Myriam Hönig**  
Leitung

Michael Seifert  
Abteilung Presse, Forschungs-  
berichterstattung, Information  
Telefon +49 7071 29-76789  
Telefax +49 7071 29-5566  
Michael.seifert@uni-tuebingen.de  
www.uni-tuebingen.de/aktuell

Wir bitten um Zusendung von  
Belegexemplaren! Danke.

Marat Minlebaev, Matthew Colonnese, Timur Tsintsadze, Anton Sirota, and Roustem Khazipov (2011). Early Gamma Oscillations Synchronize Developing Thalamus and Cortex. *Science* 334(6053):226-229

**Kontakt:**

Dr. Anton Sirota

Universität Tübingen

Werner Reichardt-Centrum für Integrative Neurowissenschaften (CIN)

Paul-Ehrlich-Straße 15

72076 Tübingen

Telefon +49 7071 29-89160

anton.sirota@cin.uni-tuebingen.de

<http://www.cin.uni-tuebingen.de/research/research-groups/physiology-of-cortico-hippocampal-interaction-sirota.php>



Anton Sirota