



Pressemitteilung



New malaria vaccine effective in clinical trial

Tübingen researchers achieve protection of up to 100 percent using fully viable malaria parasites

Tübingen, 15 February 2017

University of Tübingen researchers in collaboration with the biotech company Sanaria Inc. have demonstrated in a clinical trial that a new vaccine for malaria called Sanaria® PfSPZ-CVac has been up to 100 percent effective when assessed at 10 weeks after last dose of vaccine. For the trial, Professor Peter Kremsner and Dr. Benjamin Mordmüller of the Institute of Tropical Medicine and the German Center for Infection Research (DZIF) used malaria parasites provided by Sanaria. The vaccine incorporated fully viable – not weakened or otherwise inactivated – malaria pathogens together with the medication to combat them. Their research results have been published in the latest edition of *Nature*. DOI: 10.1038/nature21060

Malaria parasites are transmitted by the bite of female *Anopheles* mosquitoes. The *Plasmodium falciparum* parasite is responsible for most malaria infections and almost all deaths caused by the disease worldwide. Most of the previous vaccines which have been tried involved the use of individual molecules found in the pathogen. However, they were unable to provide sufficient immunity to the disease. The Tübingen study involved 67 healthy adult test persons, none of whom had previously had malaria. The best immune response was shown in a group of nine test persons who received the highest dose of the vaccine three times at four-week intervals. At the end of the trial, all nine of these individuals had 100 percent protection from the disease.

“That protection was probably caused by specific T-lymphocytes and antibody responses to the parasites in the liver,” Professor Peter Kremsner explained. The researchers analyzed the bodies’ immune reactions and identified protein patterns which will make it possible to further improve malaria vaccines, Kremsner added. The researchers injected live malaria parasites into the test subjects, at the same time preventing the development of the disease by adding chloroquine – which has been used to treat malaria for many years. This enabled the researchers to exploit the behavior of the parasites and the properties of chloroquine.

University of Tübingen
Public Relations Department

Dr. Karl Guido Rijkhoek
Director

Antje Karbe
Phone +49 7071 29-76788
+49 7071 29-76789
Fax +49 7071 29-5566
karl.rijkhoek[at]uni-tuebingen.de
antje.karbe[at]uni-tuebingen.de

www.uni-tuebingen.de/aktuell

Once the person is infected, the *Plasmodium falciparum* parasite migrates to the liver to reproduce. During its incubation period there, the human immune system could respond; but at this stage, the pathogen does not make the person sick. On top of that, chloroquine does not take effect in the liver – so it is unable to prevent the parasite from reproducing. Malaria only breaks out when the pathogen leaves the liver, entering the bloodstream and going into the red corpuscles, where it continues to reproduce and spread. As soon as the pathogen enters the bloodstream, however, it can be killed by chloroquine – and the disease cannot break out.

“By vaccinating with a live, fully active pathogen, it seems clear that we were able to set off a very strong immune response,” said study leader Benjamin Mordmüller, “Additionally, all the data we have so far indicate that what we have here is relatively stable, long-lasting protection.” In the group of test persons who demonstrated 100 percent protection after receiving a high dose three times, Mordmüller said, the protection was reliably still in place after ten weeks – and remained measurable for even longer. He added that the new vaccine showed no adverse effects on the test persons. The next step is to further test the vaccine’s effectiveness over several years in a clinical study in Gabon funded by DZIF. Malaria is one of the biggest health threats in the African nation. The University of Tübingen has worked with the Albert Schweitzer Hospital in the Gabonese town of Lambaréné and with the neighboring research institute, the Centre de Recherches Médicales de Lambaréné, for many years.

Malaria is one of the deadliest infectious diseases worldwide. The World Health Organization reports that some 214 million people became infected with malaria in the year 2015 alone. Approximately 438,000 died of the disease. Around 90 percent of those malaria deaths were in Africa. Nearly three-quarters of those who succumb to the disease are children under five. The search for a vaccine has been going on for more than a century. An effective vaccine would make it easier to control malaria; vaccination campaigns could be conducted in severely affected areas to eliminate the pathogen. Such a vaccine could also help to stop the spread of resistance to the treatment, and to better protect travelers.

Further information:

<http://www.who.int/malaria/publications/world-malaria-report-2016/report/en/>





Image: University of Tübingen



The malaria parasite *Plasmodium falciparum* under the microscope. The parasite – seen here as a dark blue ring, dot, or sickle shape – has attacked human blood cells. Photo: University of Tübingen



Images: Paul Mehnert / University of Tübingen

Publication: Benjamin Mordmüller, Güzin Surat, Heimo Lagler, Sumana Chakravarty, Andrew S. Ishizuka, Albert Lalremruata, Markus Gmeiner, Joseph J. Campo, Meral Esen, Adam J. Ruben, Jana Held, Carlos Lamsfus Calle, Juliana B. Mengue, Tamirat Gebru, Javier Ibáñez, Eric R. James, Peter F. Billingsley, Natasha KC, Anita Manoj, Tooba Murshedkar, Anusha Gunasekera, Abraham G. Eappen, Tao Li, Richard E. Stafford, MingLin Li, Phil L. Felgner, Robert A. Seder, Thomas L. Richie, B. Kim Lee Sim, Stephen L. Hoffman & Peter G. Kremsner: Sterile protection against human malaria by chemoattenuated PfSPZ vaccine, *Nature*, DOI: 10.1038/nature21060

Contact:

Professor Dr. Peter Kremsner
University Hospitals/ University of Tübingen
Institute of Tropical Medicine
Phone +49 7071 29-87179
peter.kremsner[at]uni-tuebingen.de

PD Dr. Benjamin Mordmüller
University Hospitals/ University of Tübingen
Institute of Tropical Medicine
Phone +49 7071 29-85446
benjamin.mordmueller[at]uni-tuebingen.de